

A family of multimagic squares based on large sets of orthogonal arrays

Yong Zhang¹, Kejun Chen²

1. School of Mathematical and Statistics, Yancheng Teachers University, Jiangsu 224002, China

2. Department of Mathematics, Nanjing Normal University of Special Education, Jiangsu 210038, China

3. Department of Computer Science, Lakehead University, Thunder Bay, ON P7B 5E1, Canada

Abstract

Large set of orthogonal arrays (LOA) were introduced by D. R. Stinson, and it is also used to construct multimagic squares recently. In this paper, multimagic squares based on strong double LOA are further investigated. It is proved that there exists an $\text{MS}(q^{2t-1}, t)$ for any prime power $q \geq 2t - 1$ with $t \geq 3$, which provided a new family of multimagic squares.

Keywords: Multimagic magic square, Orthogonal array, Large set.

AMS Classification: 05B15

1 Introduction

An $n \times n$ matrix A consisting of n^2 integers is a *general magic square of order n* if the sum of n elements in each row, each column and each of two diagonals are the same. The sum is the *magic number*. A general magic square of order n is a *magic square*, denoted by $\text{MS}(n)$, if the entries are n^2 consecutive integers. Usually, the (i, j) entry of a matrix A is denoted by $a_{i,j}$. A lot of work has been done on magic squares ([1, 2, 5]).

Let t be a positive integer. A general magic square M is a *general t -multimagic square* if $M^{*1}, M^{*2}, \dots, M^{*t}$ are all general magic squares, where $M^{*e} = (m_{i,j}^e)$, $e = 1, 2, \dots, t$. A general t -multimagic square of order n is a *t -multimagic square*, denoted by $\text{MS}(n, t)$, if the entries are n^2 consecutive integers. Usually, a 2-multimagic square is called a *bimagic square* and a 3-multimagic square is called a *trimagic square*.

In 2007, Derksen, Eggermont and van den Essen [6] provided a constructive procedure to make a large class of t -multimagic squares for each positive integer $t \geq 2$. There are also some new families of bimagic squares and trimagic squares, we refer the readers to [4, 9, 10]. In [9], large set of orthogonal arrays was used to construct multimagic squares, and it is also used in this paper.

An *orthogonal array of size N* , with k constraints, v levels, and strength t , denoted by $\text{OA}(N; t, k, v)$, is a $k \times N$ array with entries from a set of $v \geq 2$ symbols, having the property that in every $t \times N$ submatrix, every $t \times 1$ column vector appears the same number $\frac{N}{v^t}$ times. An $\text{OA}(N; t, k, v)$ is also denoted by $\text{OA}_\lambda(t, k, v)$, where $\lambda = \frac{N}{v^t}$. The parameter λ is the *index* of

*Corresponding to: zyyctu@gmail.com, yzhang63@lakeheadu.ca (Zhang), yctuckj@163.com(Chen)
Supported by the Natural Science Foundation of China: No.11301457(Zhang), No.11371308(Chen).

the orthogonal array. An orthogonal array is *simple* if it does not contain two identical columns. As is well known (see [5]), orthogonal arrays are of importance in design theory.

A *large set* of orthogonal arrays $\text{OA}(N; t, k, v)$, denoted by $\text{LOA}(N; t, k, v)$, is a set of v^k/N simple arrays $\text{OA}(N; t, k, v)$ such that every possible k -tuple of symbols occurs as a column in exactly one of the OAs in the set. Equivalently, the union of the OAs forms a trivial $\text{OA}(v^k; t, k, v)$. Large sets of orthogonal arrays have been used to construct resilient functions and zigzag functions by D. R. Stinson ([7, 8]). A special LOA, strong double LOA was introduced in [9] to construct t -multimagic squares.

Suppose that there is an $\text{LOA}(N; t, k, v)$, $A_s = (a_{i,j}^{(s)}), s \in I_{\frac{v^k}{N}}$. Let B_j be an array whose s -th column is the same as the j -th column of A_s , i.e.,

$$B_j = (b_{i,s}^{(j)}), b_{i,s}^{(j)} = a_{i,j}^{(s)}, i \in I_k, s \in I_{\frac{v^k}{N}}, j \in I_N,$$

Then $\{A_0, A_1, \dots, A_{\frac{v^k}{N}-1}\}$ is called a *double* $\text{LOA}(N; t, k, v)$, denoted by $\text{DLOA}(\frac{v^k}{N}, N; t, k, v)$, if $B_0, B_1, \dots, B_{N-1}$ form an $\text{LOA}(\frac{v^k}{N}; t, k, v)$. When $N = \frac{v^k}{N}$, we denote an $\text{DLOA}(\frac{v^k}{N}, N; t, k, v)$ by $\text{DLOA}(N; t, k, v)$ in short. Further, a $\text{DLOA}(N; t, k, v)$, $\{A_0, A_1, \dots, A_{N-1}\}$ is called *strong*, denoted by $\text{SDLOA}(N; t, k, v)$, if D, D' are both $\text{OA}(N; t, k, v)$ s, where D and D' are two arrays whose j -th columns are the same as j -th and $(N-1-j)$ -th column of A_j , respectively, i.e.,

$$\begin{aligned} D &= (d_{i,j})_{k \times N}, d_{i,j} = a_{i,j}^{(j)}, i \in I_k, j \in I_N, \\ D' &= (d'_{i,j})_{k \times N}, d'_{i,j} = a_{i,N-1-j}^{(j)}, i \in I_k, j \in I_N. \end{aligned}$$

We should mention that DLOA is a powerful tool to construct multimagic rectangles [11]. However, this paper mainly focus on multimagic squares. Zhang, Chen and Lei investigated the construction of multimagic squares based on SDLOA , and they proved the following.

Lemma 1.1. ([9]) *There exists an $\text{MS}(q^t, t)$ for all prime power $q \geq 2t-1$ with $t \geq 2$.*

In this paper, a further result of multimagic squares based on SDLOA is obtained as follows.

Theorem 1.2. *There exists an $\text{MS}(q^{2t-1}, t)$ for any prime power $q \geq 2t-1$ with $t \geq 3$.*

Some useful constructions of multimagic squares are reviewed in Section 2. A family of complementary t -multimagic squares are given in Section 3. The proofs of Theorem 1.2 and some useful corollaries are provided in Section 4.

2 Constructions of multimagic squares

In this section, we review some useful construction of multimagic squares presented in [6, 9, 10]. The construction of multimagic squares based on SDLOA is listed in the following.

Construction 2.1. [9] *If there exists an $\text{SDLOA}(N; t, k, v)$, then there exists an $\text{MS}(N, t)$.*

Derksen et al [6] proved the following product construction of multimagic squares.

Construction 2.2. ([6]) *If there exists an $\text{MS}(m, t)$ and there exists an $\text{MS}(n, t)$, then there exists an $\text{MS}(mn, t)$.*

Complementary t -multimagic squares was introduced in [10]. Let $I_n = \{0, 1, \dots, n-1\}$. For an $MS(n, t)$ A , the magic number of A^{*e} is denoted by $S_e(n)$, where

$$S_e(n) = \frac{\sum_{k \in I_n} k^e}{n}, e = 1, 2, \dots, t.$$

Let $B_s = (b_{i,j}^{(s)})$, $s \in I_m$, and let $B_0, B_1, \dots, B_{m-1}$ be m $MS(n, t)$ s. Then $\{B_0, B_1, \dots, B_{m-1}\}$ is called a set of complementary t -multimagic squares, denoted by m -CMS(n, t), if the following three conditions are satisfied.

$$\begin{aligned} (R1) \quad & \sum_{s \in I_m} \sum_{j \in I_n} (b_{i,j}^{(s)})^{t+1} = mS_{t+1}(n), i \in I_n; \\ (R2) \quad & \sum_{s \in I_m} \sum_{i \in I_n} (b_{i,j}^{(s)})^{t+1} = mS_{t+1}(n), j \in I_n; \\ (R3) \quad & \sum_{s \in I_m} \sum_{i \in I_n} (b_{i,i}^{(s)})^{t+1} = \sum_{s \in I_m} \sum_{i \in I_n} (b_{i,n-1-i}^{(s)})^{t+1} = mS_{t+1}(n). \end{aligned}$$

A recursive construction of t -multimagic squares were provided in [10] by using CMS.

Construction 2.3. [10] *If there is an $MS(m, t)$ and there is an m -CMS($n, t-1$), then there is an $MS(mn, t)$.*

3 Complementary t -multimagic squares based on strong double large sets of orthogonal arrays

In this section, we give a family of CMS with prime power order by making use of SDLOA.

Let F_q be a Galois field with q elements. Let $M_{k \times s}^{(t)}(F_q)$ be the set of $k \times s$ matrices over F_q with the property that any t rows are linearly independent. The following results are useful.

Lemma 3.1. [9] *If there exists a nonsingular matrix $E = (e_{i,j})_{k \times k}$ over F_q with a submatrix $E_1 \in M_{k \times s}^{(t)}(F_q)$, then there exists an LOA($q^s; t, k, q$).*

Lemma 3.2. [9] *Let $k = 2s$, $2 \leq t \leq s$. If there exists a nonsingular matrix $E = (e_{i,j})_{k \times k} = (E_1, E_2)$ over F_q such that $E_1, E_2, E_1 + E_2, E_1 - E_2$ are all in $M_{k \times s}^{(t)}(F_q)$, then there exists an SDLOA($q^s; t, k, q$).*

Let F_q^k be the set of all k -dimensional column vectors over F_q . Denote the set of a complete set of congruence classes modulo n by $\mathbb{Z}_n$, i.e., $\mathbb{Z}_n = \{\overline{0}, \overline{1}, \dots, \overline{n-1}\}$. It is known that $\mathbb{Z}_n$ is a field whenever n is prime.

Let $q = p^m$, p is prime, $n \geq 1$. A Galois field F_q can be written as a vector space over $\mathbb{Z}_p$, i.e., $F_q = \{\xi_j | \xi_j = (\overline{w}_0, \overline{w}_1, \dots, \overline{w}_{n-1})^T \in \mathbb{Z}_p^n, j = \sum_{i=0}^{m-1} w_i p^i\}$. We have $\xi_j + \xi_{q-1-j} = \xi_{q-1}$, $j \in I_q$. Denote $(\xi_{q-1}, \xi_{q-1}, \dots, \xi_{q-1})^T \in F_q^t$ by $\tilde{X}$.

We shall use SDLOA($q^t; t, 2t, q$) to construct q^t -CMS(q^t, t). For convenience, an SDLOA($q^t; t, 2t, q$) will be written as a $2t \times q^t \times q^t$ array. Further, a $2t \times q^t \times q^t$ array can be viewed as a $q^t \times q^t$ matrix C , where each element of C is a $2t$ -dimensional vector. Clearly, such a $q^t \times q^t$ matrix C is an SDLOA($q^t; t, 2t, q$) if it has the property that each row, each column,

main diagonal and back diagonal of C forms an $\text{OA}(q^t; t, 2t, q)$, and all the rows of C form an $\text{LOA}(q^t; t, 2t, q)$. We shall index the rows and the columns of C by the vectors $(x, y)^T \in F_q^t$ or just by $x + qy$. We now give a 9-CMS(9, 2) by using SDLOA.

Lemma 3.3. *There exists a 9-CMS(9, 2).*

Proof. Let $E_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix}$, $E_2 = \begin{pmatrix} 0 & 1 \\ 2 & 0 \\ 1 & 2 \\ 1 & 1 \end{pmatrix}$ and $E = (E_1, E_2)$. We can verify that E is nonsingular and $E_1, E_2, E_1 + E_2, E_1 - E_2 \in M_{4 \times 2}^{(2)}(\mathbb{Z}_3)$. We take

$$H_0 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, H_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, H_2 = \begin{pmatrix} 0 \\ 2 \end{pmatrix}, H_3 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, H_4 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, H_5 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, H_6 = \begin{pmatrix} 2 \\ 0 \end{pmatrix}, H_7 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}, H_8 = \begin{pmatrix} 2 \\ 2 \end{pmatrix};$$

$$H_0^* = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, H_1^* = \begin{pmatrix} 0 \\ 2 \end{pmatrix}, H_2^* = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, H_3^* = \begin{pmatrix} 2 \\ 1 \end{pmatrix}, H_4^* = \begin{pmatrix} 2 \\ 2 \end{pmatrix}, H_5^* = \begin{pmatrix} 2 \\ 0 \end{pmatrix}, H_6^* = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, H_7^* = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, H_8^* = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Let

$$C_i = (C_{X,Y}^{(i)}), C_{X,Y}^{(i)} = (E_1, E_2) \begin{pmatrix} X + H_i \\ Y + H_i^* \end{pmatrix} = E_1(X + H_i) + E_2(Y + H_i^*), X, Y \in \mathbb{Z}_3^2.$$

We list the $C_0, C_1, \dots, C_8$ in the following. For convenience, each entry of C_i is represented as a row vector.

$$C_0 = \begin{pmatrix} 1021 & 2012 & 0000 & 1202 & 2220 & 0211 & 1110 & 2101 & 0122 \\ 1102 & 2120 & 0111 & 1010 & 2001 & 0022 & 1221 & 2212 & 0200 \\ 1210 & 2201 & 0222 & 1121 & 2112 & 0100 & 1002 & 2020 & 0011 \\ 2000 & 0021 & 1012 & 2211 & 0202 & 1220 & 2122 & 0110 & 1101 \\ 2111 & 0102 & 1120 & 2022 & 0010 & 1001 & 2200 & 0221 & 1212 \\ 2222 & 0210 & 1201 & 2100 & 0121 & 1112 & 2011 & 0002 & 1020 \\ 0012 & 1000 & 2021 & 0220 & 1211 & 2202 & 0101 & 1122 & 2110 \\ 0120 & 1111 & 2102 & 0001 & 1022 & 2010 & 0212 & 1200 & 2221 \\ 0201 & 1222 & 2210 & 0112 & 1100 & 2121 & 0020 & 1011 & 2002 \end{pmatrix}, C_1 = \begin{pmatrix} 2120 & 0111 & 1102 & 2001 & 0022 & 1010 & 2212 & 0200 & 1221 \\ 2201 & 0222 & 1210 & 2112 & 0100 & 1121 & 2020 & 0011 & 1002 \\ 2012 & 0000 & 1021 & 2220 & 0211 & 1202 & 2101 & 0122 & 1110 \\ 0102 & 1120 & 2111 & 0010 & 1001 & 2022 & 0221 & 1212 & 2200 \\ 0210 & 1201 & 2222 & 0121 & 1112 & 2100 & 0002 & 1020 & 2011 \\ 0021 & 1012 & 2000 & 0202 & 1220 & 2211 & 0110 & 1101 & 2122 \\ 1111 & 2102 & 0120 & 1022 & 2010 & 0001 & 1200 & 2221 & 0212 \\ 1222 & 2210 & 0201 & 1100 & 2121 & 0112 & 1011 & 2002 & 0020 \\ 1000 & 2021 & 0012 & 1211 & 2202 & 0220 & 1122 & 2110 & 0101 \end{pmatrix},$$

$$C_2 = \begin{pmatrix} 0222 & 1210 & 2201 & 0100 & 1121 & 2112 & 0011 & 1002 & 2020 \\ 0000 & 1021 & 2012 & 0211 & 1202 & 2220 & 0122 & 1110 & 2101 \\ 0111 & 1102 & 2120 & 0022 & 1010 & 2001 & 0200 & 1221 & 2212 \\ 1201 & 2222 & 0210 & 1112 & 2100 & 0121 & 1020 & 2011 & 0002 \\ 1012 & 2000 & 0021 & 1220 & 2211 & 0202 & 1101 & 2122 & 0110 \\ 1120 & 2111 & 0102 & 1001 & 2022 & 0010 & 1212 & 2200 & 0221 \\ 2210 & 0201 & 1222 & 2121 & 0112 & 1100 & 2002 & 0020 & 1011 \\ 2021 & 0012 & 1000 & 2202 & 0220 & 1211 & 2110 & 0101 & 1122 \\ 2102 & 0120 & 1111 & 2010 & 0001 & 1022 & 2221 & 0212 & 1200 \end{pmatrix}, C_3 = \begin{pmatrix} 2122 & 0110 & 1101 & 2000 & 0021 & 1012 & 2211 & 0202 & 1220 \\ 2200 & 0221 & 1212 & 2111 & 0102 & 1120 & 2022 & 0010 & 1001 \\ 2011 & 0002 & 1020 & 2222 & 0210 & 1201 & 2100 & 0121 & 1112 \\ 0101 & 1122 & 2110 & 0012 & 1000 & 2021 & 0220 & 1211 & 2202 \\ 0212 & 1200 & 2221 & 0120 & 1111 & 2102 & 0001 & 1022 & 2010 \\ 0020 & 1011 & 2002 & 0201 & 1222 & 2210 & 0112 & 1100 & 2121 \\ 1110 & 2101 & 0122 & 1021 & 2012 & 0000 & 1202 & 2220 & 0211 \\ 1221 & 2212 & 0200 & 1102 & 2120 & 0111 & 1010 & 2001 & 0022 \\ 1002 & 2020 & 0011 & 1210 & 2201 & 0222 & 1121 & 2112 & 0100 \end{pmatrix},$$

$$C_4 = \begin{pmatrix} 0221 & 1212 & 2200 & 0102 & 1120 & 2111 & 0010 & 1001 & 2022 \\ 0002 & 1020 & 2011 & 0210 & 1201 & 2222 & 0121 & 1112 & 2100 \\ 0110 & 1101 & 2122 & 0021 & 1012 & 2000 & 0202 & 1220 & 2211 \\ 1200 & 2221 & 0212 & 1111 & 2102 & 0120 & 1022 & 2010 & 0001 \\ 1011 & 2002 & 0020 & 1222 & 2210 & 0201 & 1100 & 2121 & 0112 \\ 1122 & 2110 & 0101 & 1000 & 2021 & 0012 & 1211 & 2202 & 0220 \\ 2212 & 0200 & 1221 & 2120 & 0111 & 1102 & 2001 & 0022 & 1010 \\ 2020 & 0011 & 1002 & 2201 & 0222 & 1210 & 2112 & 0100 & 1121 \\ 2101 & 0122 & 1110 & 2012 & 0000 & 1021 & 2220 & 0211 & 1202 \end{pmatrix}, C_5 = \begin{pmatrix} 1020 & 2011 & 0002 & 1201 & 2222 & 0210 & 1112 & 2100 & 0121 \\ 1101 & 2122 & 0110 & 1012 & 2000 & 0021 & 1220 & 2211 & 0202 \\ 1212 & 2200 & 0221 & 1120 & 2111 & 0102 & 1001 & 2022 & 0010 \\ 2002 & 0020 & 1011 & 2210 & 0201 & 1222 & 2121 & 0112 & 1100 \\ 2110 & 0101 & 1122 & 2021 & 0012 & 1000 & 2202 & 0220 & 1211 \\ 2221 & 0212 & 1200 & 2102 & 0120 & 1111 & 2010 & 0001 & 1022 \\ 0011 & 1002 & 2020 & 0222 & 1210 & 2201 & 0100 & 1121 & 2112 \\ 0122 & 1110 & 2101 & 0000 & 1021 & 2012 & 0211 & 1202 & 2220 \\ 0200 & 1221 & 2212 & 0111 & 1102 & 2120 & 0022 & 1010 & 2001 \end{pmatrix},$$

$$C_6 = \begin{pmatrix} 0220 & 1211 & 2202 & 0101 & 1122 & 2110 & 0012 & 1000 & 2021 \\ 0001 & 1022 & 2010 & 0212 & 1200 & 2221 & 0120 & 1111 & 2102 \\ 0112 & 1100 & 2121 & 0020 & 1011 & 2002 & 0201 & 1222 & 2210 \\ 1202 & 2220 & 0211 & 1110 & 2101 & 0122 & 1021 & 2012 & 0000 \\ 1010 & 2001 & 0022 & 1221 & 2212 & 0200 & 1102 & 2120 & 0111 \\ 1121 & 2112 & 0100 & 1002 & 2020 & 0011 & 1210 & 2201 & 0222 \\ 2211 & 0202 & 1220 & 2122 & 0110 & 1101 & 2000 & 0021 & 1012 \\ 2022 & 0010 & 1001 & 2200 & 0221 & 1212 & 2111 & 0102 & 1120 \\ 2100 & 0121 & 1112 & 2011 & 0002 & 1020 & 2222 & 0210 & 1201 \end{pmatrix}, C_7 = \begin{pmatrix} 1022 & 2010 & 0001 & 1200 & 2221 & 0212 & 1111 & 2102 & 0120 \\ 1100 & 2121 & 0112 & 1011 & 2002 & 0020 & 1222 & 2210 & 0201 \\ 1211 & 2202 & 0220 & 1122 & 2110 & 0101 & 1000 & 2021 & 0012 \\ 2001 & 0022 & 1010 & 2212 & 0200 & 1221 & 2120 & 0111 & 1102 \\ 2112 & 0100 & 1121 & 2020 & 0011 & 1002 & 2201 & 0222 & 1210 \\ 2220 & 0211 & 1202 & 2101 & 0122 & 1110 & 2012 & 0000 & 1021 \\ 0010 & 1001 & 2022 & 0221 & 1212 & 2200 & 0102 & 1120 & 2111 \\ 0121 & 1112 & 2100 & 0002 & 1020 & 2011 & 0210 & 1201 & 2222 \\ 0202 & 1220 & 2211 & 0110 & 1101 & 2122 & 0021 & 1012 & 2000 \end{pmatrix},$$

$$C_8 = \begin{pmatrix} 2121 & 0112 & 1100 & 2002 & 0020 & 1011 & 2210 & 0201 & 1222 \\ 2202 & 0220 & 1211 & 2110 & 0101 & 1122 & 2021 & 0012 & 1000 \\ 2010 & 0001 & 1022 & 2221 & 0212 & 1200 & 2102 & 0120 & 1111 \\ 0100 & 1121 & 2112 & 0011 & 1002 & 2020 & 0222 & 1210 & 2201 \\ 0211 & 1202 & 2220 & 0122 & 1110 & 2101 & 0000 & 1021 & 2012 \\ 0022 & 1010 & 2001 & 0200 & 1221 & 2212 & 0111 & 1102 & 2120 \\ 1112 & 2100 & 0121 & 1020 & 2011 & 0002 & 1201 & 2222 & 0210 \\ 1220 & 2211 & 0202 & 1101 & 2122 & 0110 & 1012 & 2000 & 0021 \\ 1001 & 2022 & 0010 & 1212 & 2200 & 0221 & 1120 & 2111 & 0102 \end{pmatrix}$$

For any $i, k, l \in I_9$, if we write the (k, l) -entry of C_i in a column vector form, then each row of C_i forms a 4×9 matrix. Thus we get 81 4×9 matrix A_{ik} with the property that the l -th column of A_{ik} is the transpose of the vector of the (k, l) -entry of C_i . We list all the 4×9 matrices as follows.

$$\begin{aligned} A_{00} &= \begin{pmatrix} 120120120 \\ 000222111 \\ 210021102 \\ 120201012 \end{pmatrix}, A_{01} = \begin{pmatrix} 120120120 \\ 111000222 \\ 021102210 \\ 201012120 \end{pmatrix}, A_{02} = \begin{pmatrix} 120120120 \\ 222111000 \\ 102210021 \\ 012120201 \end{pmatrix}, \\ A_{03} &= \begin{pmatrix} 201201201 \\ 000222111 \\ 021102210 \\ 012120201 \end{pmatrix}, A_{04} = \begin{pmatrix} 201201201 \\ 111000222 \\ 102210021 \\ 120201012 \end{pmatrix}, A_{05} = \begin{pmatrix} 201201201 \\ 222111000 \\ 210021102 \\ 201012120 \end{pmatrix}, \\ A_{06} &= \begin{pmatrix} 012012012 \\ 000222111 \\ 102210021 \\ 201012120 \end{pmatrix}, A_{07} = \begin{pmatrix} 012012012 \\ 111000222 \\ 210021102 \\ 012120201 \end{pmatrix}, A_{08} = \begin{pmatrix} 012012012 \\ 222111000 \\ 021102210 \\ 120201012 \end{pmatrix}, \\ A_{10} &= \begin{pmatrix} 201201201 \\ 111000222 \\ 210021102 \\ 012120201 \end{pmatrix}, A_{11} = \begin{pmatrix} 201201201 \\ 222111000 \\ 021102210 \\ 120201012 \end{pmatrix}, A_{12} = \begin{pmatrix} 201201201 \\ 000222111 \\ 102210021 \\ 201012120 \end{pmatrix}, \\ A_{13} &= \begin{pmatrix} 012012012 \\ 111000222 \\ 021102210 \\ 201012120 \end{pmatrix}, A_{14} = \begin{pmatrix} 012012012 \\ 222111000 \\ 102210021 \\ 012120201 \end{pmatrix}, A_{15} = \begin{pmatrix} 012012012 \\ 000222111 \\ 210021102 \\ 120201012 \end{pmatrix}, \\ A_{16} &= \begin{pmatrix} 120120120 \\ 111000222 \\ 102210021 \\ 120201012 \end{pmatrix}, A_{17} = \begin{pmatrix} 120120120 \\ 222111000 \\ 210021102 \\ 201012120 \end{pmatrix}, A_{18} = \begin{pmatrix} 120120120 \\ 000222111 \\ 021102210 \\ 012120201 \end{pmatrix}, \\ A_{20} &= \begin{pmatrix} 012012012 \\ 222111000 \\ 210021102 \\ 201012120 \end{pmatrix}, A_{21} = \begin{pmatrix} 012012012 \\ 000222111 \\ 021102210 \\ 012120201 \end{pmatrix}, A_{22} = \begin{pmatrix} 012012012 \\ 111000222 \\ 102210021 \\ 120201012 \end{pmatrix}, \\ A_{23} &= \begin{pmatrix} 120120120 \\ 222111000 \\ 021102210 \\ 120201012 \end{pmatrix}, A_{24} = \begin{pmatrix} 120120120 \\ 000222111 \\ 102210021 \\ 201012120 \end{pmatrix}, A_{25} = \begin{pmatrix} 120120120 \\ 111000222 \\ 210021102 \\ 012120201 \end{pmatrix}, \\ A_{26} &= \begin{pmatrix} 201201201 \\ 222111000 \\ 102210021 \\ 012120201 \end{pmatrix}, A_{27} = \begin{pmatrix} 201201201 \\ 000222111 \\ 210021102 \\ 120201012 \end{pmatrix}, A_{28} = \begin{pmatrix} 201201201 \\ 111000222 \\ 021102210 \\ 201012120 \end{pmatrix}, \\ A_{30} &= \begin{pmatrix} 201201201 \\ 111000222 \\ 210021102 \\ 201012120 \end{pmatrix}, A_{31} = \begin{pmatrix} 201201201 \\ 222111000 \\ 021102210 \\ 012120201 \end{pmatrix}, A_{32} = \begin{pmatrix} 201201201 \\ 000222111 \\ 102210021 \\ 120201012 \end{pmatrix}, \\ A_{33} &= \begin{pmatrix} 012012012 \\ 111000222 \\ 021102210 \\ 120201012 \end{pmatrix}, A_{34} = \begin{pmatrix} 012012012 \\ 222111000 \\ 102210021 \\ 201012120 \end{pmatrix}, A_{35} = \begin{pmatrix} 012012012 \\ 000222111 \\ 210021102 \\ 012120201 \end{pmatrix}, \\ A_{36} &= \begin{pmatrix} 120120120 \\ 111000222 \\ 102210021 \\ 012120201 \end{pmatrix}, A_{37} = \begin{pmatrix} 120120120 \\ 222111000 \\ 210021102 \\ 120201012 \end{pmatrix}, A_{38} = \begin{pmatrix} 120120120 \\ 000222111 \\ 021102210 \\ 201012120 \end{pmatrix}, \\ A_{40} &= \begin{pmatrix} 012012012 \\ 222111000 \\ 210021102 \\ 120201012 \end{pmatrix}, A_{41} = \begin{pmatrix} 012012012 \\ 000222111 \\ 021102210 \\ 201012120 \end{pmatrix}, A_{42} = \begin{pmatrix} 012012012 \\ 111000222 \\ 102210021 \\ 012120201 \end{pmatrix}, \\ A_{43} &= \begin{pmatrix} 120120120 \\ 222111000 \\ 021102210 \\ 012120201 \end{pmatrix}, A_{44} = \begin{pmatrix} 120120120 \\ 000222111 \\ 102210021 \\ 120201012 \end{pmatrix}, A_{45} = \begin{pmatrix} 120120120 \\ 111000222 \\ 210021102 \\ 201012120 \end{pmatrix}, \\ A_{46} &= \begin{pmatrix} 201201201 \\ 222111000 \\ 102210021 \\ 201012120 \end{pmatrix}, A_{47} = \begin{pmatrix} 201201201 \\ 000222111 \\ 210021102 \\ 012120201 \end{pmatrix}, A_{48} = \begin{pmatrix} 201201201 \\ 111000222 \\ 021102210 \\ 120201012 \end{pmatrix}. \end{aligned}$$

$$\begin{aligned}
A_{50} &= \begin{pmatrix} 120120120 \\ 000222111 \\ 210021102 \\ 012120201 \end{pmatrix}, A_{51} = \begin{pmatrix} 120120120 \\ 111000222 \\ 021102210 \\ 120201012 \end{pmatrix}, A_{52} = \begin{pmatrix} 120120120 \\ 222111000 \\ 102210021 \\ 201012120 \end{pmatrix}, \\
A_{53} &= \begin{pmatrix} 201201201 \\ 000222111 \\ 021102210 \\ 201012120 \end{pmatrix}, A_{54} = \begin{pmatrix} 201201201 \\ 111000222 \\ 102210021 \\ 012120201 \end{pmatrix}, A_{55} = \begin{pmatrix} 201201201 \\ 222111000 \\ 210021102 \\ 120201012 \end{pmatrix}, \\
A_{56} &= \begin{pmatrix} 012012012 \\ 000222111 \\ 102210021 \\ 120201012 \end{pmatrix}, A_{57} = \begin{pmatrix} 012012012 \\ 111000222 \\ 210021102 \\ 201012120 \end{pmatrix}, A_{58} = \begin{pmatrix} 012012012 \\ 222111000 \\ 021102210 \\ 012120201 \end{pmatrix}, \\
A_{60} &= \begin{pmatrix} 012012012 \\ 222111000 \\ 210021102 \\ 012120201 \end{pmatrix}, A_{61} = \begin{pmatrix} 012012012 \\ 000222111 \\ 021102210 \\ 120201012 \end{pmatrix}, A_{62} = \begin{pmatrix} 012012012 \\ 111000222 \\ 102210021 \\ 201012120 \end{pmatrix}, \\
A_{63} &= \begin{pmatrix} 120120120 \\ 222111000 \\ 021102210 \\ 201012120 \end{pmatrix}, A_{64} = \begin{pmatrix} 120120120 \\ 000222111 \\ 102210021 \\ 012120201 \end{pmatrix}, A_{65} = \begin{pmatrix} 120120120 \\ 111000222 \\ 210021102 \\ 120201012 \end{pmatrix}, \\
A_{66} &= \begin{pmatrix} 201201201 \\ 222111000 \\ 102210021 \\ 120201012 \end{pmatrix}, A_{67} = \begin{pmatrix} 201201201 \\ 000222111 \\ 210021102 \\ 201012120 \end{pmatrix}, A_{68} = \begin{pmatrix} 201201201 \\ 111000222 \\ 021102210 \\ 012120201 \end{pmatrix}, \\
A_{70} &= \begin{pmatrix} 120120120 \\ 000222111 \\ 210021102 \\ 201012120 \end{pmatrix}, A_{71} = \begin{pmatrix} 120120120 \\ 111000222 \\ 021102210 \\ 012120201 \end{pmatrix}, A_{72} = \begin{pmatrix} 120120120 \\ 222111000 \\ 102210021 \\ 120201012 \end{pmatrix}, \\
A_{73} &= \begin{pmatrix} 201201201 \\ 000222111 \\ 021102210 \\ 120201012 \end{pmatrix}, A_{74} = \begin{pmatrix} 201201201 \\ 111000222 \\ 102210021 \\ 201012120 \end{pmatrix}, A_{75} = \begin{pmatrix} 201201201 \\ 222111000 \\ 210021102 \\ 012120201 \end{pmatrix}, \\
A_{76} &= \begin{pmatrix} 012012012 \\ 000222111 \\ 102210021 \\ 012120201 \end{pmatrix}, A_{77} = \begin{pmatrix} 012012012 \\ 111000222 \\ 210021102 \\ 120201012 \end{pmatrix}, A_{78} = \begin{pmatrix} 012012012 \\ 222111000 \\ 021102210 \\ 201012120 \end{pmatrix}, \\
A_{80} &= \begin{pmatrix} 201201201 \\ 111000222 \\ 210021102 \\ 120201012 \end{pmatrix}, A_{81} = \begin{pmatrix} 201201201 \\ 222111000 \\ 021102210 \\ 201012120 \end{pmatrix}, A_{82} = \begin{pmatrix} 201201201 \\ 000222111 \\ 102210021 \\ 012120201 \end{pmatrix}, \\
A_{83} &= \begin{pmatrix} 012012012 \\ 111000222 \\ 021102210 \\ 012120201 \end{pmatrix}, A_{84} = \begin{pmatrix} 012012012 \\ 222111000 \\ 102210021 \\ 120201012 \end{pmatrix}, A_{85} = \begin{pmatrix} 012012012 \\ 000222111 \\ 210021102 \\ 201012120 \end{pmatrix}, \\
A_{86} &= \begin{pmatrix} 120120120 \\ 111000222 \\ 102210021 \\ 201012120 \end{pmatrix}, A_{87} = \begin{pmatrix} 120120120 \\ 222111000 \\ 210021102 \\ 012120201 \end{pmatrix}, A_{88} = \begin{pmatrix} 120120120 \\ 000222111 \\ 021102210 \\ 120201012 \end{pmatrix}.
\end{aligned}$$

For any $i \in I_9$, $X + H_i$ and $Y + H_i^*$ both run over $\mathbb{Z}_3^2$ when X and Y both run over $\mathbb{Z}_3^2$. Since E is nonsingular and $E_1, E_2, E_1 + E_2, E_1 - E_2 \in M_{4 \times 2}^{(2)}(\mathbb{Z}_3)$, $\{A_{i0}, A_{i1}, \dots, A_{i8}\}$ is an SDLOA(2, 4, 3) by the proof of Lemma 3.2. Thus C_i forms an SDLOA(9; 2, 4, 3) $\{A_{i0}, A_{i1}, \dots, A_{i8}\}$ for each $i \in I_9$. One can check that the 9 SDLOA(2, 4, 3)s have the following properties:

- (1) for fixed $k \in I_9$, the $A_{0k}, A_{1k}, \dots, A_{8k}$ forms a LOA(2, 4, 3);
- (2) for fixed $l \in I_9$, the l -th column of all $A_{ik} (i, k \in I_9)$ forms a LOA(2, 4, 3);
- (3) all the k -th column of $A_{ik}, i, k \in I_9$ forms a LOA(2, 4, 3), and the $(8 - k)$ -th column of $A_{ik}, i, k \in I_9$ forms a LOA(2, 4, 3).

For any C_i , replacing the arbitrary (k, l) -entry (u, v, x, y) of C_i by $u + 3v + 9x + 27y$, we get a matrix $\hat{C}_i$. By the proof of Construction 2.1, $\hat{C}_0, \hat{C}_1, \dots, \hat{C}_8$ are MS(9, 2)s. We list $\hat{C}_0, \hat{C}_1, \dots, \hat{C}_8$ as follows.

$$\begin{aligned}
\hat{C}_0 &= \begin{pmatrix} 46 & 65 & 0 & 61 & 26 & 42 & 13 & 32 & 75 \\ 58 & 23 & 39 & 10 & 29 & 72 & 52 & 71 & 6 \\ 16 & 35 & 78 & 49 & 68 & 3 & 55 & 20 & 36 \\ 2 & 45 & 64 & 44 & 60 & 25 & 77 & 12 & 31 \\ 41 & 57 & 22 & 74 & 9 & 28 & 8 & 51 & 70 \\ 80 & 15 & 34 & 5 & 48 & 67 & 38 & 54 & 19 \\ 63 & 1 & 47 & 24 & 43 & 62 & 30 & 76 & 14 \\ 21 & 40 & 59 & 27 & 73 & 11 & 69 & 7 & 53 \\ 33 & 79 & 17 & 66 & 4 & 50 & 18 & 37 & 56 \end{pmatrix}, \hat{C}_1 = \begin{pmatrix} 23 & 39 & 58 & 29 & 72 & 10 & 71 & 6 & 52 \\ 35 & 78 & 16 & 68 & 3 & 49 & 20 & 36 & 55 \\ 65 & 0 & 46 & 26 & 42 & 61 & 32 & 75 & 13 \\ 57 & 22 & 41 & 9 & 28 & 74 & 51 & 70 & 8 \\ 15 & 34 & 80 & 48 & 67 & 5 & 54 & 19 & 38 \\ 45 & 64 & 2 & 60 & 25 & 44 & 12 & 31 & 77 \\ 40 & 59 & 21 & 73 & 11 & 27 & 7 & 53 & 69 \\ 79 & 17 & 33 & 4 & 50 & 66 & 37 & 56 & 18 \\ 1 & 47 & 63 & 43 & 62 & 24 & 76 & 14 & 30 \end{pmatrix}, \hat{C}_2 = \begin{pmatrix} 78 & 16 & 35 & 3 & 49 & 68 & 36 & 55 & 20 \\ 0 & 46 & 65 & 42 & 61 & 26 & 75 & 13 & 32 \\ 39 & 58 & 23 & 72 & 10 & 29 & 6 & 52 & 71 \\ 34 & 80 & 15 & 67 & 5 & 48 & 19 & 38 & 54 \\ 64 & 2 & 45 & 25 & 44 & 60 & 31 & 77 & 12 \\ 22 & 41 & 57 & 28 & 74 & 9 & 70 & 8 & 51 \\ 17 & 33 & 79 & 50 & 66 & 4 & 56 & 18 & 37 \\ 47 & 63 & 1 & 62 & 24 & 43 & 14 & 30 & 76 \\ 59 & 21 & 40 & 11 & 27 & 73 & 53 & 69 & 7 \end{pmatrix}, \\
\hat{C}_3 &= \begin{pmatrix} 77 & 12 & 31 & 2 & 45 & 64 & 44 & 60 & 25 \\ 8 & 51 & 70 & 41 & 57 & 22 & 74 & 9 & 28 \\ 38 & 54 & 19 & 80 & 15 & 34 & 5 & 48 & 67 \\ 30 & 76 & 14 & 63 & 1 & 47 & 24 & 43 & 62 \\ 69 & 7 & 53 & 21 & 40 & 59 & 27 & 73 & 11 \\ 18 & 37 & 56 & 33 & 79 & 17 & 66 & 4 & 50 \\ 13 & 32 & 75 & 46 & 65 & 0 & 61 & 26 & 42 \\ 52 & 71 & 6 & 58 & 23 & 39 & 10 & 29 & 72 \\ 55 & 20 & 36 & 16 & 35 & 78 & 49 & 68 & 3 \end{pmatrix}, \hat{C}_4 = \begin{pmatrix} 51 & 70 & 8 & 57 & 22 & 41 & 9 & 28 & 74 \\ 54 & 19 & 38 & 15 & 34 & 80 & 48 & 67 & 5 \\ 12 & 31 & 77 & 45 & 64 & 2 & 60 & 25 & 44 \\ 7 & 53 & 69 & 40 & 59 & 21 & 73 & 11 & 27 \\ 37 & 56 & 18 & 79 & 17 & 33 & 4 & 50 & 66 \\ 76 & 14 & 30 & 1 & 47 & 63 & 43 & 62 & 24 \\ 71 & 6 & 52 & 23 & 39 & 58 & 29 & 72 & 10 \\ 20 & 36 & 55 & 35 & 78 & 16 & 68 & 3 & 49 \\ 32 & 75 & 13 & 65 & 0 & 46 & 26 & 42 & 61 \end{pmatrix}, \hat{C}_5 = \begin{pmatrix} 19 & 38 & 54 & 34 & 80 & 15 & 67 & 5 & 48 \\ 31 & 77 & 12 & 64 & 2 & 45 & 25 & 44 & 60 \\ 70 & 8 & 51 & 22 & 41 & 57 & 28 & 74 & 9 \\ 56 & 18 & 37 & 17 & 33 & 79 & 50 & 66 & 4 \\ 14 & 30 & 76 & 47 & 63 & 1 & 62 & 24 & 43 \\ 53 & 69 & 7 & 59 & 21 & 40 & 11 & 27 & 73 \\ 36 & 55 & 20 & 78 & 16 & 35 & 3 & 49 & 68 \\ 75 & 13 & 32 & 0 & 46 & 65 & 42 & 61 & 26 \\ 6 & 52 & 71 & 39 & 58 & 23 & 72 & 10 & 29 \end{pmatrix}, \\
\hat{C}_6 &= \begin{pmatrix} 24 & 43 & 62 & 30 & 76 & 14 & 63 & 1 & 47 \\ 27 & 73 & 11 & 69 & 7 & 53 & 21 & 40 & 59 \\ 66 & 4 & 50 & 18 & 37 & 56 & 33 & 79 & 17 \\ 61 & 26 & 42 & 13 & 32 & 75 & 46 & 65 & 0 \\ 10 & 29 & 72 & 52 & 71 & 6 & 58 & 23 & 39 \\ 49 & 68 & 3 & 55 & 20 & 36 & 16 & 35 & 78 \\ 44 & 60 & 25 & 77 & 12 & 31 & 2 & 45 & 64 \\ 74 & 9 & 28 & 8 & 51 & 70 & 41 & 57 & 22 \\ 5 & 48 & 67 & 38 & 54 & 19 & 80 & 15 & 34 \end{pmatrix}, \hat{C}_7 = \begin{pmatrix} 73 & 11 & 27 & 7 & 53 & 69 & 40 & 59 & 21 \\ 4 & 50 & 66 & 37 & 56 & 18 & 79 & 17 & 33 \\ 43 & 62 & 24 & 76 & 14 & 30 & 1 & 47 & 63 \\ 29 & 72 & 10 & 71 & 6 & 52 & 23 & 39 & 58 \\ 68 & 3 & 49 & 20 & 36 & 55 & 35 & 78 & 16 \\ 26 & 42 & 61 & 32 & 75 & 13 & 65 & 0 & 46 \\ 9 & 28 & 74 & 51 & 70 & 8 & 57 & 22 & 41 \\ 48 & 67 & 5 & 54 & 19 & 38 & 15 & 34 & 80 \\ 60 & 25 & 44 & 12 & 31 & 77 & 45 & 64 & 2 \end{pmatrix}, \hat{C}_8 = \begin{pmatrix} 50 & 66 & 4 & 56 & 18 & 37 & 17 & 33 & 79 \\ 62 & 24 & 43 & 14 & 30 & 76 & 47 & 63 & 1 \\ 11 & 27 & 73 & 53 & 69 & 7 & 59 & 21 & 40 \\ 3 & 49 & 68 & 36 & 55 & 20 & 78 & 16 & 35 \\ 42 & 61 & 26 & 75 & 13 & 32 & 0 & 46 & 65 \\ 72 & 10 & 29 & 6 & 52 & 71 & 39 & 58 & 23 \\ 67 & 5 & 48 & 19 & 38 & 54 & 34 & 80 & 15 \\ 25 & 44 & 60 & 31 & 77 & 12 & 64 & 2 & 45 \\ 28 & 74 & 9 & 70 & 8 & 51 & 22 & 41 & 57 \end{pmatrix}.
\end{aligned}$$

By the property (1), for fixed $k \in I_9$, the union of the columns of $A_{0k}, A_{1k}, \dots, A_{8k}$ is exactly the set F_3^4 . So the numbers of all the k -th rows of $\hat{C}_0, \hat{C}_1, \dots, \hat{C}_8$ forms the set I_{81} exactly, which is independent of the choice of k . So the condition R_1 in the definition of CMS is satisfied. In the same way, by the properties (2) and (3) the conditions R_2 and R_3 are satisfied, respectively. Thus $\{\hat{C}_0, \hat{C}_1, \dots, \hat{C}_8\}$ is a 9-CMS(9, 2). $\square$

We now give a sufficient condition of q^t -CMS(q^t, t) based on LOA while keeping the idea of 9-CMS(9, 2) in mind.

Lemma 3.4. *If there exist q^t SDLOA($q^t; t, 2t, q$)s over F_q , $C_H = (C_{X,Y}^{(H)})$, $X, Y, H \in F_q^t$, such that*

$$\begin{aligned}
R_X &= \{C_{X,Y}^{(H)} | Y, H \in F_q^t\}, X \in F_q^t, \\
T_Y &= \{C_{X,Y}^{(H)} | X, H \in F_q^t\}, Y \in F_q^t, \\
U &= \{C_{X,X}^{(H)} | X, H \in F_q^t\}, \\
U' &= \{C_{X,\tilde{X}-X}^{(H)} | X, H \in F_q^t\}
\end{aligned}$$

are $2q^t + 2$ LOA($q^t; t, 2t, q$)s, then there exists a q^t -CMS(q^t, t).

Proof. Let $\hat{C}_H = (\hat{C}_{X,Y}^{(H)})$, where

$$\hat{C}_{X,Y}^{(H)} = j_0 + j_1 q + \dots + j_{2t-1} q^{2t-1} = (1, q, \dots, q^{2t-1})(j_0, j_1, \dots, j_{2t-1})^T$$

if $C_{X,Y}^{(H)} = (\xi_{j_0}, \xi_{j_1}, \dots, \xi_{j_{2t-1}})^T$, $X, Y, H \in F_q^t$. By the proof of Lemma 2.1, $\hat{C}_H, H \in F_q^t$ are q^t MS(q^t, t)s. We shall prove that $\{\hat{C}_H | H \in F_q^t\}$ is a q^t -CMS(q^t, t).

In fact, for fixed $X \in F_q^t$, since R_X forms an LOA($q^t; t, 2t, q$), we have

$$\{\hat{C}_{X,Y}^{(H)} | Y, H \in F_q^t\} = I_{q^{2t}}.$$

Thus $\sum_{H \in F_q^t} \sum_{Y \in F_q^t} (\hat{C}_{X,Y}^{(H)})^{t+1} = \sum_{d=0}^{q^{2t}-1} d^{t+1}, X \in F_q^t$. But $\sum_{d=0}^{q^{2t}-1} d^{t+1} = q^t S_{t+1}(q^t)$. So we have

$$\sum_{H \in F_q^t} \sum_{Y \in F_q^t} (\hat{C}_{X,Y}^{(H)})^{t+1} = q^t S_{t+1}(q^t), X \in F_q^t.$$

Similarly, for fixed Y , T_Y forms an $\text{LOA}(q^t; t, 2t, q)$. So we have

$$\sum_{H \in F_q^t} \sum_{X \in F_q^t} (\hat{C}_{X,Y}^{(H)})^{t+1} = q^t S_{t+1}(q^t), Y \in F_q^t;$$

In the same way, since U and U' both forms $\text{LOA}(q^t; t, 2t, q)$ s. So

$$\sum_{H \in F_q^t} \sum_{X \in F_q^t} (\hat{C}_{X,X}^{(H)})^{t+1} = \sum_{H \in F_q^t} \sum_{X \in F_q^t} (\hat{C}_{X, \tilde{X}-X}^{(H)})^{t+1} = q^t S_{t+1}(q^t).$$

So, $\{\hat{C}_H | H \in F_q^t\}$ is a q^t -CMS(q^t, t). □

Now the technique of matrix is used to construct a q^t -CMS(q^t, t).

Lemma 3.5. *Let $t \geq 2$ and $d \in F_q$. If there exists a matrix $E = (e_{i,j})_{2t \times 2t} = (E_1, E_2)$ over F_q such that*

- (1) $E_1, E_2, E_1 + E_2, E_1 - E_2$ are all in $M_{2t \times t}^{(t)}(F_q)$,
 - (2) $E, (E_1, E_1 + dE_2), (E_2, E_1 + dE_2), (E_1 + E_2, E_1 + dE_2), (E_1 - E_2, E_1 + dE_2)$ are nonsingular,
- then there exists a q^t -CMS(q^t, t).*

Proof. Let $C = (C_{X,Y})$, where $C_{X,Y} = (E_1, E_2) \begin{pmatrix} X \\ Y \end{pmatrix}, X, Y \in F_q^t$. By the proof of Lemma 3.2, C forms an $\text{SDLOA}(q^t; t, 2t, q)$ over F_q . We shall construct a q^t -CMS(q^t, t) by making use of the coordinate translations of C .

Let $H \in F_q^t$ and $H^* = dH$. Let

$$C_H = (C_{X,Y}^{(H)}), H \in F_q^t,$$

where

$$C_{X,Y}^{(H)} = C_{X+H, Y+H^*} = (E_1, E_2) \begin{pmatrix} X+H \\ Y+H^* \end{pmatrix}, X, Y \in F_q^t.$$

i.e.,

$$C_{X,Y}^{(H)} = E_1 X + E_2 Y + (E_1 + dE_2)H, X, Y \in F_q^t.$$

Let $\hat{C}_H$ be the same as defined in Lemma 3.4. We shall show that $\{\hat{C}_H | H \in F_q^t\}$ is a q^t -CMS(q^t, t).

For fixed $H \in F_q^t$, let $K_H = (E_1, E_2) \begin{pmatrix} H \\ H^* \end{pmatrix}$, then

$$C_{X,Y}^{(H)} = (E_1, E_2) \begin{pmatrix} X \\ Y \end{pmatrix} + K_H, X, Y \in F_q^t,$$

which means that C_H is obtained by adding K_H to each column of the 3-dimensional array C . Since C is an $\text{SDLOA}(q^t; t, 2t, q)$, it is obvious that C_H is also an $\text{SDLOA}(q^t; t, 2t, q)$. Let $R_X(X \in F_q^t), T_Y(Y \in F_q^t), U, U'$ be the same as in Lemma 3.4. By Lemma 3.4, it remains to prove that $R_X(X \in F_q^t), T_Y(Y \in F_q^t), U, U'$ are all $\text{LOA}(q^t; t, 2t, q)$ s. We have

$$\begin{aligned} R_X &= \{E_1X + E_2Y + (E_1 + dE_2)H | Y, H \in F_q^t\} \\ &= \{E_1X + (E_2, E_1 + dE_2) \begin{pmatrix} Y \\ H \end{pmatrix} | Y, H \in F_q^t\}, \quad X \in F_q^t \end{aligned}$$

Since $E_2 \in M_{2t \times t}^{(t)}(F_q)$ and $(E_2, E_1 + dE_2)$ is nonsingular, R_X forms an $\text{LOA}(q^t; t, 2t, q)$ for each $X \in F_q^t$. Similarly, since $E_1 \in M_{2t \times t}^{(t)}(F_q)$ and $(E_1, E_1 + dE_2)$ is nonsingular, T_Y forms an $\text{LOA}(q^t; t, 2t, q)$ for each $Y \in F_q^t$.

Now we consider U, U' , by Lemma 3.1 we have

$$\begin{aligned} U &= \{E_1X + E_2X + (E_1 + dE_2)H | X, H \in F_q^t\} \\ &= \{(E_1 + E_2, E_1 + dE_2) \begin{pmatrix} X \\ H \end{pmatrix} | X, H \in F_q^t\}. \end{aligned}$$

and

$$\begin{aligned} U' &= \{E_1X + E_2(\tilde{X} - X) + (E_1 + dE_2)H | X, H \in F_q^t\} \\ &= \{(E_1 - E_2, E_1 + dE_2) \begin{pmatrix} X \\ H \end{pmatrix} + E_2\tilde{X} | X, H \in F_q^t\} \end{aligned}$$

since $E_1 + E_2, E_1 - E_2 \in M_{2t \times t}^{(t)}(F_q)$ and $(E_1 + E_2, E_1 + dE_2), (E_1 - E_2, E_1 + dE_2)$ are nonsingular, U, U' are both $\text{LOA}(q^t; t, 2t, q)$ s. The proof is completed. $\square$

Lemma 3.6. *Let q be a prime power and $q \geq 4$. If $M_{2t \times t}^{(t)}(F_q) \neq \emptyset$, then there exists a q^t -CMS(q^t, t).*

Proof. Let x be a primitive element in F_q . Since $q \geq 4$, which indicates that $x, x^2 \notin \{0, 1, -1\}$ and $x \neq x^2$. Suppose that E_{11}, E_{21} are two $t \times t$ matrices such that $E_1 = \begin{pmatrix} E_{11} \\ E_{21} \end{pmatrix} \in M_{2t \times t}^{(t)}(F_q)$. Let

$$E = (E_1, E_2) = \begin{pmatrix} E_{11} & E_{12} \\ E_{21} & E_{22} \end{pmatrix} = \begin{pmatrix} E_{11} & xE_{11} \\ E_{21} & x^2E_{21} \end{pmatrix},$$

and $d = x$. It is readily checked that E satisfies the conditions in Lemma 3.5. Therefore there exists a q^t -CMS(q^t, t) by Lemma 3.5. $\square$

Theorem 3.7. *There exists a q^t -CMS(q^t, t) for any prime power $q \geq 2t - 1$ with $t \geq 2$.*

Proof. For $q = 3$ and $t = 2$, there exist 9-CMS(9, 2) by Lemma 3.3. For $q \geq \max\{2t - 1, 4\}$ and $t \geq 2$, let x be a primitive element of F_q and

$$E = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 1 \\ 1 & x & x^2 & \cdots & x^{t-1} \\ 1 & x^2 & x^4 & \cdots & x^{2(t-1)} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & x^{2t-2} & x^{2(2t-2)} & \cdots & x^{(t-1)(2t-2)} \end{pmatrix}.$$

It is readily checked that any t rows of E are linearly independent, so $E \in M_{2t \times t}^{(t)}(F_q)$. Thus, there exists a q^t -CMS(q^t, t) by Lemma 3.6. $\square$

4 Main results

Theorem 1.2 There exists an MS(q^{2t-1}, t) for any prime power $q \geq 2t - 1$ with $t \geq 3$.

Proof. For any prime power $q \geq 2t - 1$ with $t \geq 3$, there exists an MS(q^t, t) by Lemma 1.1 and exists an q^{t-1} -CMS($q^{t-1}, t - 1$) by Theorem 3.7. So there exists an MS(q^{2t-1}, t) by Construction 2.3. $\square$

Remark The MS($q^{2t-1}, 2t - 1$)s coming from Theorem 1.2 with prime power $q < 20$ are listed as follows.

$$\begin{array}{cccccccc}
 (5^5, 3) & (7^5, 3) & (8^5, 3) & (9^5, 3) & (11^5, 3) & (13^5, 3) & (16^5, 3) & (17^5, 3) & (19^5, 3) \\
 (7^7, 4) & (8^7, 4) & (9^7, 4) & (11^7, 4) & (13^7, 4) & (16^7, 4) & (17^7, 4) & (19^7, 4) & \\
 & (9^9, 5) & (11^9, 5) & (13^9, 5) & (16^9, 5) & (17^9, 5) & (19^9, 5) & & \\
 & & (11^{11}, 6) & (13^{11}, 6) & (16^{11}, 6) & (17^{11}, 6) & (19^{11}, 6) & & \\
 & & & (13^{13}, 7) & (16^{13}, 7) & (17^{13}, 7) & (19^{13}, 7) & & \\
 & & & & (16^{15}, 8) & (17^{15}, 8) & (19^{15}, 8) & & \\
 & & & & & (17^{17}, 9) & (19^{17}, 9) & &
 \end{array}$$

We should point out that those marked in red can't be obtained from Lemma 1.1 directly. Further, we have the following new family of multimagic squares.

Corollary 4.1. *There exists an MS(q^{2t-1}, t) for any prime power $2t - 1 \leq q < 4t - 3$ with $t \geq 3$.*

Proof. For a $(2t - 1)$ -multimagic square, Lemma 1.1 provides an MS($q^{2(2t-1)-1}, 2t - 1$) for any prime power $q \geq 2(2t - 1) - 1$ with $2t - 1 \geq 2$, i.e., an MS($q^{4t-3}, 2t - 1$) for any prime power $q \geq 4t - 3$ with $t \geq 2$. But Theorem 1.2 provides an MS($q^{2t-1}, 2t - 1$) for any prime power $q \geq 2t - 1$ with $t \geq 3$. So the MS(q^{2t-1}, t)s for prime power $2t - 1 \leq q < 4t - 3$ with $t \geq 3$ are covered by Theorem 1.2 but are not covered by Lemma 1.1. $\square$

For fixed q and fixed t , those multimagic squares come from Lemma 1.1 and Theorem 1.2 are both single case, but the following provided a family ones.

Corollary 4.2. *There exists an MS(q^m, t) for any prime power q with $q \geq 4t - 5$ and $m \geq t \geq 3$.*

Proof. Let $t \geq 3$ and let $m = t + k, k \geq 0$. For $0 \leq k \leq t - 2$ and $q \geq 2(t + k) - 1$, by Lemma 1.1 there exists an MS($q^{t+k}, t + k$) and hence there exists an MS(q^{t+k}, t). For $k = t - 1$, there exists an MS(q^{t+k}, t) when $q \geq 2t - 1$ by Theorem 1.2. For any $m \geq 2t$, we can write $m = \sum_{i=0}^{t-1} (t + i)n_i$, where $n_0, n_1, \dots, n_{t-1}$ are nonnegative integers. By Construction 2.2, we get an MS(q^m, t). The proof is completed. $\square$

Open problem Find more MS(q^t, t) and MS(q^{2t-1}, t) for prime power $q < 2t - 1$ with $t \geq 3$.

References

- [1] G. Abe, Unsolved problems on magic squares, Discrete Math. 127 (1994) 3-13.
- [2] W. S. Andrews, Magic Squares and Cubes, 2nd ed., Dover, New York, 1960.
- [3] C. Boyer, Multimagic squares, www.multimagic.com.

- [4] K. Chen, W. Li, Existence of normal bimagic squares, *Discrete Math.* 312(2012) 3077-3086.
- [5] C. J. Colbourn, J. H. Dinitz (eds.), *Handbook of combinatorial designs*, 2nd Edition, Chapman and Hall/CRC, Boca Raton FL, 2007.
- [6] H. Derksen, C. Eggermont, A. van den Essen, Multimagic squares, *Amer. Math. Monthly* 114 (2007) 703-713.
- [7] D. R. Stinson, Resilient functions and large sets of orthogonal arrays, *Congr. Numer.* 92(1993) 105-110.
- [8] D. R. Stinson, Some results on nonlinear zigzag functions, *J. Combin. Math. Combin. Comput.* 29 (1999) 127-138.
- [9] Y. Zhang, K. Chen, J. Lei, Large sets of orthogonal arrays and multimagic squares, *J. Combin. Des.* 21(2013) 390-403.
- [10] Y. Zhang, K. Chen, W. Li, A generalization of product construction of multimagic squares, *Advances in Mathematics (China)*, to appear. (arXiv:1512.07287 [math.CO])
- [11] Y. Zhang, J. Lei, Multimagic rectangles based on large sets of orthogonal arrays, *Discrete Math.* 313(18)(2013): 1823-1831.